10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HNRPK antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11741

Specification

HNRPK antibody - N-terminal region - Product Information

Application	**IHC, WB**
Primary Accession	P61978
Other Accession	NM_031262, NP_112552
Reactivity	**Human, Mouse, Rat, Rabbit, Zebrafish, Goat, Horse, Bovine, Dog**
Predicted	**Human, Mouse, Rat, Rabbit, Zebrafish, Chicken, Bovine**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**51kDa KDa**

HNRPK antibody - N-terminal region - Additional Information

Gene ID 3190

Alias Symbol	**CSBP, TUNP, HNRPK**

Other Names
Heterogeneous nuclear ribonucleoprotein K, hnRNP K, Transformation up-regulated nuclear protein, TUNP, HNRNPK, HNRPK

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-HNRPK antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
HNRPK antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HNRPK antibody - N-terminal region - Protein Information

Name HNRNPK

Synonyms HNRPK

Function
One of the major pre-mRNA-binding proteins. Binds tenaciously to poly(C) sequences. Likely to play a role in the nuclear metabolism of hnRNAs, particularly for pre-mRNAs that contain cytidine-rich sequences. Can also bind poly(C) single-stranded DNA. Plays an important role in p53/TP53 response to DNA damage, acting at the level of both transcription activation and

repression. When sumoylated, acts as a transcriptional coactivator of p53/TP53, playing a role in p21/CDKN1A and 14-3-3 sigma/SFN induction (By similarity). As far as transcription repression is concerned, acts by interacting with long intergenic RNA p21 (lincRNA-p21), a non-coding RNA induced by p53/TP53. This interaction is necessary for the induction of apoptosis, but not cell cycle arrest. As part of a ribonucleoprotein complex composed at least of ZNF827, HNRNPL and the circular RNA circZNF827 that nucleates the complex on chromatin, may negatively regulate the transcription of genes involved in neuronal differentiation (PubMed:33174841).


Cellular Location
Cytoplasm. Nucleus, nucleoplasm. Cell projection, podosome. Note=Recruited to p53/TP53-responsive promoters, in the presence of functional p53/TP53 (PubMed:16360036). In case of ASFV infection, there is a shift in the localization which becomes predominantly nuclear (PubMed:18775702)

HNRPK antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HNRPK antibody - N-terminal region - Images



Rabbit Anti-HNRPK Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



WB Suggested Anti-HNRPK Antibody Titration: 0.2-1 µg/ml
ELISA Titer: 1:62500
Positive Control: HepG2 cell lysate

HNRPK antibody - N-terminal region - References

Nakagawa,J., (2006) Br. J. Cancer 94 (4), 586-592Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.