

LSM6 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI11787**Specification**

LSM6 antibody - middle region - Product Information

Application	WB
Primary Accession	P62312
Other Accession	NM_007080 , NP_009011
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Yeast, Bovine, Dog
Predicted	Human, Mouse, Rat, Zebrafish, Chicken, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	9kDa KDa

LSM6 antibody - middle region - Additional Information**Gene ID** 11157

Alias Symbol	YDR378°C
Other Names	
U6 snRNA-associated Sm-like protein LSm6, LSM6	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-LSM6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

LSM6 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

LSM6 antibody - middle region - Protein Information**Name** LSM6**Function**

Plays a role in pre-mRNA splicing as component of the U4/U6- U5 tri-snRNP complex that is involved in spliceosome assembly, and as component of the precatalytic spliceosome (spliceosome B complex) (PubMed:<<http://www.uniprot.org/citations/28781166>>28781166). The heptameric LSM2-8 complex binds specifically to the 3'-terminal U-tract of U6 snRNA (PubMed:<<http://www.uniprot.org/citations/10523320>>10523320). Component of LSm protein complexes, which are involved in RNA processing and may function in a chaperone-like manner, facilitating the efficient association

of RNA processing factors with their substrates. Component of the cytoplasmic LSM1-LSM7 complex, which is thought to be involved in mRNA degradation by activating the decapping step in the 5'-to-3' mRNA decay pathway (Probable).


Cellular Location

Cytoplasm. Nucleus

LSM6 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LSM6 antibody - middle region - Images

WB Suggested Anti-LSM6 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: HCT15 cell lysate

LSM6 is supported by BioGPS gene expression data to be expressed in HCT15