

NXF3 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11809**Specification**

NXF3 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	O9H4D5
Other Accession	NM_022052 , NP_071335
Reactivity	Human, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Rabbit, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 58kDa KDa

NXF3 antibody - C-terminal region - Additional Information**Gene ID** 56000**Other Names**

Nuclear RNA export factor 3, TAP-like protein 3, TAPL-3, NXF3, TAPL3

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-NXF3 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

NXF3 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NXF3 antibody - C-terminal region - Protein Information**Name** NXF3**Synonyms** TAPL3**Function**

May function as a tissue-specific nuclear mRNA export factor.

Cellular Location

Nucleus. Cytoplasm. Note=Shuttles between the nucleus and the cytoplasm

Tissue Location

Expressed at high level in testis and at low level in a small number of tissues

NXF3 antibody - C-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NXF3 antibody - C-terminal region - Images



Rabbit Anti-NXF3 Antibody
Paraffin Embedded Tissue: Human Heart
Cellular Data: Myocardial cells
Antibody Concentration: 4.0-8.0 $\mu\text{g/ml}$
Magnification: 400X



Rabbit Anti-NXF3 Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 $\mu\text{g/ml}$
Magnification: 400X



WB Suggested Anti-NXF3 Antibody Titration: 1.25µg/ml

Positive Control: RPMI 8226 cell lysate

NXF3 is supported by BioGPS gene expression data to be expressed in RPMI 8226

NXF3 antibody - C-terminal region - References

Jun,L., (2001) Curr. Biol. 11 (18), 1381-1391
Reconstitution and Storage: For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.