

NOC4L antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11835**Specification**

NOC4L antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q9BVI4
Other Accession	NM_024078 , NP_076983
Reactivity	Human, Mouse, Rat, Pig, Horse, Bovine
Predicted	Human, Mouse, Rat, Pig, Horse, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57kDa KDa

NOC4L antibody - C-terminal region - Additional Information**Gene ID** 79050**Alias Symbol** **NOC4, NET49, UTP19****Other Names**

Nucleolar complex protein 4 homolog, NOC4 protein homolog, NOC4-like protein, Nucleolar complex-associated protein 4-like protein, NOC4L

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NOC4L antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

NOC4L antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NOC4L antibody - C-terminal region - Protein Information**Name** NOC4L**Cellular Location**


Nucleus membrane; Multi-pass membrane protein. Nucleus, nucleolus

NOC4L antibody - C-terminal region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NOC4L antibody - C-terminal region - Images



Rabbit Anti-NOC4L Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



WB Suggested Anti-NOC4L Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate

NOC4L antibody - C-terminal region - References

Rual, J.F., (2005) Nature 437 (7062), 1173-1178
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.