

ADAD2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11846**Specification**

ADAD2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8NCV1
Other Accession	NM_139174 , NP_631913
Reactivity	Human, Rat, Rabbit, Horse, Bovine, Dog
Predicted	Human, Rabbit, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71kDa KDa

ADAD2 antibody - C-terminal region - Additional Information**Gene ID** 161931**Alias Symbol** **FLJ00337, TENRL****Other Names**

Adenosine deaminase domain-containing protein 2, Testis nuclear RNA-binding protein-like, ADAD2, TENRL

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ADAD2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ADAD2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ADAD2 antibody - C-terminal region - Protein Information**Name** ADAD2 ([HGNC:30714](#))**Synonyms** TENRL**Function**

Required for male fertility and normal male germ cell differentiation.

Cellular Location


Nucleus {ECO:0000250|UniProtKB:Q9D5P4}. Cytoplasm {ECO:0000250|UniProtKB:Q9D5P4}. Note=Diffusely cytoplasmic early in pachytene spermatocytes and coalesces into several perinuclear granules by late pachynema. {ECO:0000250|UniProtKB:Q9D5P4}

ADAD2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADAD2 antibody - C-terminal region - Images



WB Suggested Anti-ADAD2 Antibody Titration: 0.2-1 µg/ml
Positive Control: HepG2 cell lysate

ADAD2 antibody - C-terminal region - References

Ota,T., (2004) Nat. Genet. 36 (1), 40-45 Reconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.