10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NKD2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11870

Specification

NKD2 antibody - N-terminal region - Product Information

Application	**WB**
Primary Accession	Q96EK8
Other Accession	NM_033120, AAH12176
Reactivity	**Human, Mouse, Rat, Horse**
Predicted	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**34kDa KDa**

NKD2 antibody - N-terminal region - Additional Information

Alias Symbol	**Naked2**

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-NKD2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
NKD2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NKD2 antibody - N-terminal region - Protein Information


NKD2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

NKD2 antibody - N-terminal region - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



WB Suggested Anti-NKD2 Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate

NKD2 antibody - N-terminal region - References

Strausberg,R. Direct SubmissionReconstitution and Storage:For short term use, store at 2-8C up to 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.