

ADAMTS4 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11908**Specification**

ADAMTS4 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q8NEK2
Other Accession	AAH30812
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Pig, Horse, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 37kDa KDa

ADAMTS4 antibody - N-terminal region - Additional Information

Alias Symbol **ADAMTS-2, ADAMTS-4, ADMP-1, KIAA0688**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ADAMTS4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ADAMTS4 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ADAMTS4 antibody - N-terminal region - Protein Information**ADAMTS4 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ADAMTS4 antibody - N-terminal region - Images



Rabbit Anti-ADAMTS4 Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



WB Suggested Anti-ADAMTS4 Antibody Titration: 0.2-1 µg/ml
Positive Control: HepG2 cell lysate