

AGER antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11920**Specification**

AGER antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	A2BF17
Other Accession	NM_172197 , NP_751947
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

AGER antibody - C-terminal region - Additional Information

Alias Symbol	MGC22357, RAGE
Target/Specificity	
Isoform 2 specific	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-AGER antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

AGER antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


AGER antibody - C-terminal region - Protein Information**AGER antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AGER antibody - C-terminal region - Images

Rabbit Anti-AGER Antibody
Paraffin Embedded Tissue: Human Lung
Cellular Data: Alveolar cells
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



70 kDa
60 kDa
48 kDa
36 kDa
21 kDa

WB Suggested Anti-AGER Antibody Titration: 0.2-1 µg/ml
Positive Control: HepG2 cell lysate