10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

AGER antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11921

Specification

AGER antibody - C-terminal region - Product Information

Application	**IHC, WB**
Primary Accession	A2BFI7
Other Accession	NM_172197, NP_751947
Reactivity	**Human**
Predicted	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**38kDa KDa**

AGER antibody - C-terminal region - Additional Information

Alias Symbol	**MGC22357, RAGE**

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 100 ul of distilled water. Final anti-AGER antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
AGER antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

AGER antibody - C-terminal region - Protein Information


AGER antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

AGER antibody - C-terminal region - Images


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Rabbit Anti-AGER Antibody
Paraffin Embedded Tissue: Human Kidney
Cellular Data: Epithelial cells of renal tubule
Antibody Concentration: 4.0-8.0 µg/ml
Magnification: 400X



WB Suggested Anti-AGER Antibody Titration: 2.5µg/ml
Positive Control: Jurkat cell lysate
AGER is supported by BioGPS gene expression data to be expressed in Jurkat



WB Suggested Anti-AGER antibody Titration: 1 µg/ml
Sample Type: Human K562