

F2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11936**Specification**

F2 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	P18292
Other Accession	NM_022924 , NP_075213
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Rat, Sheep, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 70kDa KDa

F2 antibody - N-terminal region - Additional Information

Alias Symbol	F2
Other Names	
Prothrombin, 3.4.21.5, Coagulation factor II, Activation peptide fragment 1, Activation peptide fragment 2, Thrombin light chain, Thrombin heavy chain, F2	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-F2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

F2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

F2 antibody - N-terminal region - Protein Information**Name** F2**Function**


Thrombin, which cleaves bonds after Arg and Lys, converts fibrinogen to fibrin and activates factors V, VII, VIII, XIII, and, in complex with thrombomodulin, protein C. Functions in blood homeostasis, inflammation and wound healing.

F2 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F2 antibody - N-terminal region - Images



WB Suggested Anti-F2 Antibody Titration: 1.0 µg/ml
Positive Control: Rat Lung