

FGA antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI11937**Specification**

FGA antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q4QQH7
Other Accession	NM_021871 , NP_068657
Reactivity	Human, Mouse, Rat, Pig, Goat, Sheep, Horse, Bovine, Dog
Predicted Host	Human, Mouse, Horse, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 71kDa KDa

FGA antibody - N-terminal region - Additional Information

Alias Symbol **Fib2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-FGA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


FGA antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

FGA antibody - N-terminal region - Protein Information**FGA antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FGA antibody - N-terminal region - Images



Rabbit Anti-FGA Antibody
Paraffin Embedded Tissue: Human Skin
Cellular Data: Squamous epithelial cells
Antibody Concentration: 4.0-8.0 $\mu\text{g/ml}$
Magnification: 400X



WB Suggested Anti-FGA Antibody Titration: 5.0 $\mu\text{g/ml}$
Positive Control: Jurkat cell lysate