

NUDT9 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12027**Specification**

NUDT9 antibody - C-terminal region - Product Information

Application	IHC, WB
Primary Accession	Q8NG26
Other Accession	NM_198038 , NP_932155
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Sheep, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33kDa kDa

NUDT9 antibody - C-terminal region - Additional Information

Alias Symbol **NUDT10**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-NUDT9 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


NUDT9 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

NUDT9 antibody - C-terminal region - Protein Information**NUDT9 antibody - C-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NUDT9 antibody - C-terminal region - Images



Human Skin



WB Suggested Anti-NUDT9 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:62500

Positive Control: Jurkat cell lysate

NUDT9 antibody - C-terminal region - References

Clark, H.F., et al., (2003) Genome Res. 13(10), 2265-2270. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.