10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Fkhl18 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12048

Specification

Fkhl18 antibody - C-terminal region - Product Information

Application	**WB**
Primary Accession	Q61574
Other Accession	NM_010226, NP_034356
Reactivity	**Mouse, Rat, Horse, Bovine, Guinea Pig**
Predicted	**Mouse, Rat, Pig, Horse, Bovine, Guinea Pig**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**35kDa KDa**

Fkhl18 antibody - C-terminal region - Additional Information

Gene ID 14239

Alias Symbol	**FREAC10, Fkh3, Foxs1, Fkhl18**

Other Names
Forkhead box protein S1, Forkhead-like 18 protein, Forkhead-related transcription factor 10, FREAC-10, Transcription factor FKH-3, Foxs1, Fkh3, Fkhl18, Freac10

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-Fkhl18 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
Fkhl18 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Fkhl18 antibody - C-terminal region - Protein Information

Name Foxs1

Synonyms Fkh3, Fkhl18, Freac10

Function
Transcriptional repressor that suppresses transcription from the FASLG, FOXO3 and FOXO4 promoters. May have a role in the organization of the testicular vasculature.

Cellular Location
Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089, ECO:0000269|PubMed:18288644}


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Fkhl18 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Fkhl18 antibody - C-terminal region - Images



WB Suggested Anti-Fkhl18 Antibody Titration: 0.2-1 µg/ml
Positive Control: Mouse Brain