

DMRT2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12088**Specification**

DMRT2 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	O9NQR6
Other Accession	NM_181872 , NP_870987
Reactivity	Human, Rabbit
Predicted	Human, Rabbit
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62kDa KDa

DMRT2 antibody - N-terminal region - Additional Information**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-DMRT2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


DMRT2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DMRT2 antibody - N-terminal region - Protein Information**DMRT2 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DMRT2 antibody - N-terminal region - Images



Human kidney



WB Suggested Anti-DMRT2 Antibody Titration: 1.25µg/ml
Positive Control: HepG2 cell lysate

DMRT2 antibody - N-terminal region - References

Ottolenghi, C., (2000) Genomics 64(2), 179-186
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.