

TBC1D16 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12113**Specification**

TBC1D16 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q8TBP0
Other Accession	NM_019020 , NP_061893
Reactivity	Human, Mouse, Rat, Zebrafish, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Zebrafish, Pig, Chicken, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	86kDa KDa

TBC1D16 antibody - N-terminal region - Additional Information**Gene ID** 125058

Alias Symbol	FLJ20748, MGC25062
Other Names	
TBC1 domain family member 16, TBC1D16	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TBC1D16 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TBC1D16 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TBC1D16 antibody - N-terminal region - Protein Information**Name** TBC1D16**Function**


May act as a GTPase-activating protein for Rab family protein(s).

TBC1D16 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TBC1D16 antibody - N-terminal region - Images



WB Suggested Anti-TBC1D16 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: HT1080 cell lysate