

DNASE2B antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12124**Specification**

DNASE2B antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5VXD0
Other Accession	NM_058248 , NP_490649
Reactivity	Human, Mouse, Rat, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Rat, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17kDa KDa

DNASE2B antibody - C-terminal region - Additional Information

Alias Symbol **DLAD**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-DNASE2B antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


DNASE2B antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DNASE2B antibody - C-terminal region - Protein Information**DNASE2B antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DNASE2B antibody - C-terminal region - Images



WB Suggested Anti-DNASE2B Antibody Titration: 1.25µg/ml
Positive Control: Jurkat cell lysate

DNASE2B antibody - C-terminal region - References

MacLea, K.S., (2003) Gene 305(1), 1-12 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.