10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HERC6 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12243

Specification

HERC6 antibody - N-terminal region - Product Information

Application	**WB**
Primary Accession	Q8IVU3
Other Accession	NM_017912, NP_060382
Reactivity	**Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog**
Predicted	**Human, Rabbit, Horse, Bovine, Guinea Pig, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**115kDa KDa**

HERC6 antibody - N-terminal region - Additional Information

Gene ID 55008

Alias Symbol	**FLJ20637**

Other Names
Probable E3 ubiquitin-protein ligase HERC6, 6.3.2.-, HECT domain and RCC1-like domain-containing protein 6, HERC6

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-HERC6 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
HERC6 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

HERC6 antibody - N-terminal region - Protein Information

Name HERC6

Function
E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates.

Cellular Location
Cytoplasm, cytosol.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999


Tissue Location
Detected in brain, heart, placenta and testis.

HERC6 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HERC6 antibody - N-terminal region - Images



WB Suggested Anti-HERC6 Antibody Titration: 0.2-1 µg/ml
Positive Control: ACHN cell lysate