

ITCH antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12267**Specification**

ITCH antibody - N-terminal region - Product Information

Application	WB
Primary Accession	A9Z1X3
Other Accession	NM_031483 , NP_113671
Reactivity	Human, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Rabbit, Horse, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 99kDa KDa

ITCH antibody - N-terminal region - Additional Information

Alias Symbol **AIF4, AIP4, NAPP1, dj468O1.1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ITCH antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ITCH antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ITCH antibody - N-terminal region - Protein Information**ITCH antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ITCH antibody - N-terminal region - Images



WB Suggested Anti-ITCH Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: COLO205 cell lysate