

TMEM176A antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12291**Specification**

TMEM176A antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q96HP8
Other Accession	NM_018487 , NP_060957
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	26kDa KDa

TMEM176A antibody - N-terminal region - Additional Information**Gene ID** 55365**Alias Symbol** **GS188, HCA112****Other Names**

Transmembrane protein 176A, Hepatocellular carcinoma-associated antigen 112, TMEM176A, HCA112

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-TMEM176A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TMEM176A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

TMEM176A antibody - N-terminal region - Protein Information**Name** TMEM176A**Synonyms** HCA112**Cellular Location**


Membrane; Multi-pass membrane protein

TMEM176A antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TMEM176A antibody - N-terminal region - Images



WB Suggested Anti-TMEM176A Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: THP-1 cell lysate