

SLC22A16 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12372**Specification**

SLC22A16 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	O96RU0
Other Accession	NM_033125 , NP_149116
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	63kDa KDa

SLC22A16 antibody - N-terminal region - Additional Information

Alias Symbol **CT2, FLIPT2, OCT6, OKB1, dj261K5.1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-SLC22A16 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions


SLC22A16 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SLC22A16 antibody - N-terminal region - Protein Information**SLC22A16 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC22A16 antibody - N-terminal region - Images



Human kidney



WB Suggested Anti-SLC22A16 Antibody Titration: 2.5 µg/ml
Positive Control: HepG2 cell lysate

SLC22A16 antibody - N-terminal region - References

Okabe, M., (2005) Biochem. Biophys. Res. Commun. 333(3), 754-762 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.