

POR antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12410**Specification**

POR antibody - middle region - Product Information

Application	IHC, WB
Primary Accession	Q63HL4
Other Accession	NM_000941 , NP_000932
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Zebrafish, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	77kDa KDa

POR antibody - middle region - Additional Information

Alias Symbol	CPR, CYPOR, DKFZp686G04235, FLJ26468, P450R
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-POR antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


POR antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

POR antibody - middle region - Protein Information**POR antibody - middle region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

POR antibody - middle region - Images



0



WB Suggested Anti-POR Antibody Titration: 0.25µg/ml
Positive Control: 293T cells lysate

POR antibody - middle region - References

Gomes, L.G., (er) J. Clin. Endocrinol. Metab. (2008) In press Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. Publications: Lee, J.W., Kim, Y.H., Yoon, S. & Lee, S.K. Cytochrome P450 system expression and DNA adduct formation in the liver of *Zacco platypus* following waterborne Benzo(a)pyrene exposure: implications for biomarker determination. Environ. Toxicol. (2012). doi:10.1002/tox.21833 WB, Bovine, Human, Dog, Rabbit, Guinea pig, Rat, Mouse, Horse, Zebrafish 23192953