10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Ppic Antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12412

Specification

Ppic Antibody - C-terminal region - Product Information

Application	**WB**
Primary Accession	P30412
Other Accession	NM_008908, NP_032934
Reactivity	**Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog**
Predicted	**Human, Mouse, Rat, Pig, Bovine, Dog**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**23kDa KDa**

Ppic Antibody - C-terminal region - Additional Information

Gene ID 19038

Alias Symbol	**CyP-20°C**

Other Names
Peptidyl-prolyl cis-trans isomerase C, PPIase C, 5.2.1.8, Cyclophilin C, Rotamase C, Ppic, Cypc

Format
Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage
Add 50 ul of distilled water. Final anti-Ppic antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions
Ppic Antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Ppic Antibody - C-terminal region - Protein Information

Name Ppic {ECO:0000312|MGI:MGI:97751}

Function
PPIase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding.

Cellular Location
Cytoplasm.

Tissue Location
Specifically expressed in kidney.


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Ppic Antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Ppic Antibody - C-terminal region - Images



WB Suggested Anti-Ppic Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Heart