

EDA antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12437**Specification**

EDA antibody - middle region - Product Information

Application	WB
Primary Accession	B7ZLU4
Other Accession	NM_001005609 , NP_001005609
Reactivity	Human, Mouse, Rat, Rabbit, Goat, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41kDa KDa

EDA antibody - middle region - Additional Information

Alias Symbol	ED1, ED1-A1, ED1-A2, EDA1, EDA2, HED, XHED, XLHED, ODT1, STHAGX1
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-EDA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


EDA antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

EDA antibody - middle region - Protein Information**EDA antibody - middle region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EDA antibody - middle region - Images



WB Suggested Anti-EDA Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: HepG2 cell lysate

EDA antibody - middle region - References

Zhao, J., (2008) Br. J. Dermatol. 158(3), 614-617 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.