

WDR33 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12439**Specification**

WDR33 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O9NUL1
Other Accession	NM_001006622 , NP_001006623
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Zebrafish, Pig, Chicken, Horse, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

WDR33 antibody - N-terminal region - Additional Information

Alias Symbol **FLJ11294, WDC146, NET14**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-WDR33 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


WDR33 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

WDR33 antibody - N-terminal region - Protein Information**WDR33 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WDR33 antibody - N-terminal region - Images



WB Suggested Anti-WDR33 Antibody Titration: 0.2-1 μ g/ml
Positive Control: THP-1 cell lysate