

MFAP3L antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12450**Specification**

MFAP3L antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O75121
Other Accession	NM_021647 , NP_067679
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Rabbit, Pig, Chicken, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45kDa KDa

MFAP3L antibody - N-terminal region - Additional Information**Gene ID** 9848**Alias Symbol** NYD-sp9**Other Names**

Microfibrillar-associated protein 3-like, Testis development protein NYD-SP9, MFAP3L, KIAA0626

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-MFAP3L antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

MFAP3L antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

MFAP3L antibody - N-terminal region - Protein Information**Name** MFAP3L**Synonyms** KIAA0626**Function**

May participate in the nuclear signaling of EGFR and MAPK1/ERK2. May have a role in metastasis.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm Note=Mainly localized

in the nucleus (PubMed:24735981)


Tissue Location

Highly expressed in testis. {ECO:0000269|Ref.1}.

MFAP3L antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MFAP3L antibody - N-terminal region - Images

WB Suggested Anti-MFAP3L Antibody Titration: 2.5µg/ml
Positive Control: Jurkat cell lysate

MFAP3L antibody - N-terminal region - References

Yang, C.B., Unpublished (2003) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.