

C3orf17 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12494**Specification**

C3orf17 antibody - middle region - Product Information

Application	WB
Primary Accession	Q6NW34
Other Accession	NM_015412 , NP_056227
Reactivity	Human, Dog
Predicted	Human, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64kDa KDa

C3orf17 antibody - middle region - Additional Information**Gene ID** 25871**Alias Symbol** DKFZP434F2021, NET17**Other Names**

Uncharacterized protein C3orf17, C3orf17

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C3orf17 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C3orf17 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

C3orf17 antibody - middle region - Protein Information**Name** NEPRO ([HGNC:24496](#))**Synonyms** C3orf17**Function**

May play a role in cortex development as part of the Notch signaling pathway. Downstream of Notch may repress the expression of proneural genes and inhibit neuronal differentiation thereby maintaining neural progenitors. May also play a role in preimplantation embryo development.

Cellular Location


Nucleus {ECO:0000250|UniProtKB:Q8R2U2}. Nucleus, nucleolus {ECO:0000250|UniProtKB:Q8R2U2}

C3orf17 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C3orf17 antibody - middle region - Images



WB Suggested Anti-C3orf17 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: 721_B cell lysate

C3orf17 antibody - middle region - References

Nousiainen, M., (2006) Proc. Natl. Acad. Sci. U.S.A. 103(14), 5391-5396 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.