

USP48 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12500**Specification**

USP48 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q86UV5-7
Other Accession	NM_001032730 , AAH67261
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70kDa KDa

USP48 antibody - C-terminal region - Additional Information

Alias Symbol	DKFZp762M1713, MGC132556, MGC14879, RAP1GA1, USP31
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-USP48 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


USP48 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

USP48 antibody - C-terminal region - Protein Information**USP48 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

USP48 antibody - C-terminal region - Images



WB Suggested Anti-USP48 Antibody Titration: 0.2-1 µg/ml

Positive Control: HepG2 cell lysate

USP48 is strongly supported by BioGPS gene expression data to be expressed in Human HepG2 cells