

TCP10L antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI12634**Specification**

TCP10L antibody - middle region - Product Information

Application	IHC, WB
Primary Accession	Q8TDR4
Other Accession	NM_144659 , NP_653260
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24kDa KDa

TCP10L antibody - middle region - Additional Information**Gene ID** 140290**Alias Symbol** PRED77, TCP10A-2, C21orf77**Other Names**

T-complex protein 10A homolog 2, T-complex protein 10A-2, TCP10A-2, TCP10-like, TCP10L

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-TCP10L antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

TCP10L antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

TCP10L antibody - middle region - Protein Information**Name** TCP10L**Function**

May be involved in transcriptional regulation. Has in vitro transcription inhibition activity. Acts as a tumor suppressor in hepatocellular carcinoma (HCC) cells.

Cellular Location

Nucleus

Tissue Location


Expressed in liver and testis. Expressed in the seminiferous tubules (at protein level).

TCP10L antibody - middle region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TCP10L antibody - middle region - Images



Human Heart



WB Suggested Anti-TCP10L Antibody Titration: 1.25µg/ml
Positive Control: HepG2 cell lysate