

DDT antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12649**Specification**

DDT antibody - N-terminal region - Product Information

Application	WB
Primary Accession	B5MCS6
Other Accession	NM_001355 , NP_001346
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig
Predicted Host	Human, Mouse, Pig
Clonality	Rabbit
Calculated MW	Polyclonal 13kDa KDa

DDT antibody - N-terminal region - Additional Information

Alias Symbol **DDCT**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-DDT antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


DDT antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DDT antibody - N-terminal region - Protein Information**DDT antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDT antibody - N-terminal region - Images



WB Suggested Anti-DDT Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:312500

Positive Control: Human Liver

DDT antibody - N-terminal region - References

Ewing, R.M., Mol. Syst. Biol. 3, 89 (2007) Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.