

DDOST antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12696**Specification**

DDOST antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q5VWA5
Other Accession	NM_005216 , NP_005207
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Sheep, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Zebrafish, Pig, Chicken, Horse, Bovine, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46kDa KDa

DDOST antibody - N-terminal region - Additional Information

Alias Symbol	AGE-R1, KIAA0115, MGC2191, OK/SW-cl.45, OST, OST48, WBP1, AGER1, OKSWcl45
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-DDOST antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

DDOST antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

DDOST antibody - N-terminal region - Protein Information**DDOST antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DDOST antibody - N-terminal region - Images

WB Suggested Anti-DDOST Antibody Titration: 5.0µg/ml
Positive Control: Jurkat cell lysate

DDOST antibody - N-terminal region - References

Shibatani, T., (2005) Biochemistry 44(16), 5982-5992 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.