

MLF2 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12697**Specification**

MLF2 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q15773
Other Accession	NM_005439 , NP_005430
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Mouse, Rat, Chicken, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 28kDa KDa

MLF2 antibody - C-terminal region - Additional Information**Gene ID** 8079

Alias Symbol	NTN4
Other Names	
Myeloid leukemia factor 2, Myelodysplasia-myeloid leukemia factor 2, MLF2	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-MLF2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

MLF2 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

MLF2 antibody - C-terminal region - Protein Information**Name** MLF2**Cellular Location**

Cytoplasm. Nucleus.

Tissue Location


Ubiquitously expressed.

MLF2 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MLF2 antibody - C-terminal region - Images



WB Suggested Anti-MLF2 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate

MLF2 antibody - C-terminal region - References

Walker, L.C., (et al) Breast Cancer Res. Treat. (2007) In press Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.