

RTN2 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12700**Specification**

RTN2 antibody - N-terminal region - Product Information

Application	IHC, WB
Primary Accession	A8K7F2
Other Accession	NM_206900 , NP_996783
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51kDa KDa

RTN2 antibody - N-terminal region - Additional Information

Alias Symbol **NSP2, NSPL1**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-RTN2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


RTN2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

RTN2 antibody - N-terminal region - Protein Information**RTN2 antibody - N-terminal region - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RTN2 antibody - N-terminal region - Images



Human Liver



WB Suggested Antibody Titration: 2.5 µg/ml

Positive Control: Jurkat

RTN2 antibody - N-terminal region - References

Oertle, T., (2003) FASEBJ. 17(10), 1238-1247. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.