

Vamp7 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12813**Specification**

Vamp7 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	P70280
Other Accession	NM_011515 , NP_035645
Reactivity	Human, Mouse, Rat, Rabbit, Zebrafish, Sheep, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Chicken, Sheep, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25kDa kDa

Vamp7 antibody - C-terminal region - Additional Information**Gene ID** 20955**Alias Symbol** **Sybl1, TI-VAMP, VAMP-7****Other Names**

Vesicle-associated membrane protein 7, VAMP-7, Synaptobrevin-like protein 1, Vamp7, Sybl1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Vamp7 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Vamp7 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Vamp7 antibody - C-terminal region - Protein Information**Name** Vamp7**Synonyms** Sybl1**Function**

Involved in the targeting and/or fusion of transport vesicles to their target membrane during transport of proteins from the early endosome to the lysosome. Required for heterotypic fusion of late endosomes with lysosomes and homotypic lysosomal fusion. Required for calcium regulated lysosomal exocytosis. Involved in the export of chylomicrons from the endoplasmic reticulum to the cis Golgi. Required for exocytosis of mediators during eosinophil and neutrophil degranulation,

and target cell killing by natural killer cells. Required for focal exocytosis of late endocytic vesicles during phagosome formation.

Cellular Location


Cytoplasmic vesicle, secretory vesicle membrane; Single-pass type IV membrane protein. Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrane protein. Late endosome membrane; Single-pass type IV membrane protein. Lysosome membrane; Single-pass type IV membrane protein. Endoplasmic reticulum membrane; Single-pass type IV membrane protein. Cytoplasmic vesicle, phagosome membrane; Single-pass type IV membrane protein. Synapse, synaptosome Note=In immature neurons expression is localized in vesicular structures in axons and dendrites while in mature neurons it is localized to the somatodendritic region. Colocalizes with LAMP1 in kidney cells. Localization to the endoplasmic reticulum membrane was observed in the intestine but not in liver or kidney (By similarity)

Vamp7 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Vamp7 antibody - C-terminal region - Images



WB Suggested Anti-Vamp7 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse liver