

ZNF138 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12816**Specification**

ZNF138 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	P52744
Other Accession	NM_006524 , NP_006515
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31kDa KDa

ZNF138 antibody - N-terminal region - Additional Information**Gene ID** 7697

Alias Symbol	pHZ-32
Other Names	
Zinc finger protein 138, ZNF138	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-ZNF138 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

ZNF138 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF138 antibody - N-terminal region - Protein Information**Name** ZNF138**Function**

May be involved in transcriptional regulation as a repressor.

Cellular Location


Nucleus.

ZNF138 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF138 antibody - N-terminal region - Images



WB Suggested Anti-ZNF138 Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:62500

Positive Control: 293T cell lysate

ZNF138 antibody - N-terminal region - References

Tommerup, N. (1995) Genomics 27(2), 259-264. Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.