

**RUNDC2A antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI12830****Specification**

---

**RUNDC2A antibody - N-terminal region - Product Information**

|                   |                                                          |
|-------------------|----------------------------------------------------------|
| Application       | IHC, WB                                                  |
| Primary Accession | <a href="#">O9HA26</a>                                   |
| Other Accession   | <a href="#">NM_032167</a> , <a href="#">NP_115543</a>    |
| Reactivity        | Human, Mouse, Rat, Rabbit, Zebrafish, Horse, Bovine, Dog |
| Predicted Host    | Human, Bovine                                            |
| Clonality         | Rabbit                                                   |
| Calculated MW     | Polyclonal<br>42kDa KDa                                  |

**RUNDC2A antibody - N-terminal region - Additional Information**

Alias Symbol **FLJ12363**

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 100 ul of distilled water. Final anti-RUNDC2A antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

RUNDC2A antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


**RUNDC2A antibody - N-terminal region - Protein Information****RUNDC2A antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**RUNDC2A antibody - N-terminal region - Images**

---



Human Intestine



WB Suggested Anti-RUNDC2A Antibody Titration: 5.0µg/ml  
Positive Control: HepG2 cell lysate

### **RUNDC2A antibody - N-terminal region - References**

Bonaldo, M.F., (1996) Genome Res. 6(9), 791-806 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.