

hCG_20426 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12847**Specification**

hCG_20426 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	E5RJM6
Other Accession	XM_001126121 , XP_001126121
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42kDa KDa

hCG_20426 antibody - C-terminal region - Additional Information**Gene ID** 441869**Alias Symbol** **LOC441869, hCG_20426****Other Names**

Ankyrin repeat domain-containing protein 65, ANKRD65

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-hCG_20426 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

hCG_20426 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


hCG_20426 antibody - C-terminal region - Protein Information**Name** ANKRD65**hCG_20426 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

hCG_20426 antibody - C-terminal region - Images



WB Suggested Anti-hCG_20426 Antibody Titration: 0.2-1 μ g/ml
Positive Control: Jurkat cell lysate