

IL28RA antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12873**Specification**

IL28RA antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q5VTX8
Other Accession	NM_173064 , NP_775087
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54kDa KDa

IL28RA antibody - N-terminal region - Additional Information

Alias Symbol	CRF2/12, IFNLR, IFNLR1, LICR2, il-28r1, IL-28R1, IL28RA
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-IL28RA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


IL28RA antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

IL28RA antibody - N-terminal region - Protein Information**IL28RA antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL28RA antibody - N-terminal region - Images



WB Suggested Anti-IL28RA Antibody Titration: 0.2-1 µg/ml

ELISA Titer: 1:312500

Positive Control: Jurkat cell lysate

IL28RA antibody - N-terminal region - References

Chae, S.C., (2006) Exp. Mol. Med. 38(3), 302-309 Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. Publications: Mucha, J., Majchrzak, K., Taciak, B., Hellmén, E. & Król, M. MDSCs mediate angiogenesis and predispose cancer in mammary tumor cells for metastasis via IL-28/IL-28RA (IFN-λ) signaling. PLoS One 9, e103249 (2014). WB, Human, Guinea pig, Dog, Rat, Bovine, Horse, Rabbit, Mouse 25075523