

PRDX2 antibody - middle region

Rabbit Polyclonal Antibody Catalog # Al12896

Specification

PRDX2 antibody - middle region - Product Information

Application Primary Accession Other Accession Reactivity

Predicted

Host Clonality Calculated MW IHC, WB P35704 NM_005809, NP_005800 Human, Mouse, Rat, Rabbit, Zebrafish, Goat, Sheep, Horse, Yeast, Bovine, Guinea Pig, Dog Human, Mouse, Rat, Zebrafish, Pig, Sheep, Bovine, Guinea Pig Rabbit Polyclonal 22kDa KDa

PRDX2 antibody - middle region - Additional Information

Gene ID 29338

Alias Symbol

MGC4104, NKEFB, PRP, PRX2, PRXII, TDPX1, TSA, TPX1

Other Names

Peroxiredoxin-2, 1.11.1.15, Thiol-specific antioxidant protein, TSA, Thioredoxin peroxidase 1, Thioredoxin-dependent peroxide reductase 1, Prdx2, Tdpx1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-PRDX2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions PRDX2 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

PRDX2 antibody - middle region - Protein Information

Name Prdx2

Synonyms Tdpx1

Function

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative



stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2).

Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:P32119}.

PRDX2 antibody - middle region - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PRDX2 antibody - middle region - Images



Liver



Rabbit Anti-PRDX2 Antibody Formalin Fixed Paraffin Embedded Tissue: Human Pineal Tissue Observed Staining: Cytoplasmic in cell bodies and processes of pinealocytes

Primary Antibody


Concentration: 1:100 Other Working Concentrations: 1/600 Secondary Antibody: Donkey anti-Rabbit-Cy3 Secondary Antibody

Concentration: 1:200

Magnification: 20X

Exposure Time: 0.5 - 2.0 sec





WB Suggested Anti-PRDX2 Antibody Titration: 0.2-1 µg/ml ELISA Titer: 1:312500 Positive Control: Hela cell lysate PRDX2 is supported by BioGPS gene expression data to be expressed in HeLa

PRDX2 antibody - middle region - References

Fang, J., (2007) Proc. Natl. Acad. Sci. U.S.A. 104 (47), 18742-18747 Reconstitution and Storage: Forshortter muse, storeat 2-8 Cupto1 week. For long terms to rage, store at -20 Cinsmallaliquots to prevent freeze - thaw cup science.