

FNTA antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12913**Specification**

FNTA antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A6NJW0
Other Accession	NM_001018677 , NP_001018197
Reactivity	Human, Rat, Rabbit, Pig, Sheep, Horse, Bovine, Dog
Predicted Host	Human, Rat, Sheep, Bovine, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 36kDa KDa

FNTA antibody - C-terminal region - Additional Information

Alias Symbol **FPTA, MGC99680, PGGT1A, PTAR2**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-FNTA antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


FNTA antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

FNTA antibody - C-terminal region - Protein Information**FNTA antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FNTA antibody - C-terminal region - Images



WB Suggested Anti-FNTA Antibody Titration: 1.25 μ g/ml
Positive Control: HepG2 cell lysate

FNTA antibody - C-terminal region - References

Veluthakal, R., (2007) Diabetes 56(1), 204-210 Reconstitution and Storage: For short term use, store at 2-8C upto 1 week. For long term storage, store at -20C in small aliquots to prevent freeze-thaw cycles.