

CDY1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12925**Specification**

CDY1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	A8K8F3
Other Accession	NM_004680 , NP_004671
Reactivity	Human, Mouse, Rat, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human
Clonality	Rabbit
Calculated MW	Polyclonal 62kDa KDa

CDY1 antibody - C-terminal region - Additional Information

Alias Symbol **CDY, CDY1A**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-CDY1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


CDY1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

CDY1 antibody - C-terminal region - Protein Information**CDY1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDY1 antibody - C-terminal region - Images



WB Suggested Anti-CDY1 Antibody Titration: 2.5µg/ml
Positive Control: HepG2 cell lysate

CDY1 antibody - C-terminal region - References

Stouffs, K., (2008) Hum. Reprod. 23(5), 1193-1199
Reconstitution and Storage: For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.