

Mettl4 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI12957**Specification**

Mettl4 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	D3ZD59
Other Accession	NM_001191814 , NP_001178743
Reactivity	Human, Mouse, Rat, Rabbit, Sheep, Horse, Bovine, Guinea Pig, Dog
Predicted	Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53kDa KDa

Mettl4 antibody - C-terminal region - Additional Information

Alias Symbol	RGD1306451, Mettl4, HsT661, AV296509, 2410198H06Rik, A730091E08Rik
--------------	--

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Mettl4 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.


Precautions

Mettl4 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Mettl4 antibody - C-terminal region - Protein Information**Mettl4 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mettl4 antibody - C-terminal region - Images

WB Suggested Anti-Mettl4 Antibody Titration: 0.2-1 μ g/ml

ELISA Titer: 1:12500

Positive Control: Rat Liver