

Zik1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13166**Specification**

Zik1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q80YP6
Other Accession	NM_009577 , NP_033603
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Mouse
Clonality	Rabbit
Calculated MW	Polyclonal 53kDa KDa

Zik1 antibody - C-terminal region - Additional Information**Gene ID** 22775**Other Names**

Zinc finger protein interacting with ribonucleoprotein K, Zik1

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Zik1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Zik1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Zik1 antibody - C-terminal region - Protein Information**Name** Zik1**Function**

May be a transcriptional repressor.

Cellular Location

Nucleus.

Tissue Location


Expressed in ovary and liver, and at lower levels in brain and muscle.

Zik1 antibody - C-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Zik1 antibody - C-terminal region - Images



WB Suggested Anti-Zik1 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Kidney

Zik1 antibody - C-terminal region - References

Denisenko O.N., et al. J. Biol. Chem. 271:27701-27706(1996).
Carninci P., et al. Science 309:1559-1563(2005).