

ING1 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13184**Specification**

ING1 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q5T9H0
Other Accession	NM_198219 , NP_937862
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32kDa KDa

ING1 antibody - C-terminal region - Additional Information

Alias Symbol	p24ING1c, p33, p33ING1, p33ING1b, p47, p47ING1a
--------------	---

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 100 ul of distilled water. Final anti-ING1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


ING1 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

ING1 antibody - C-terminal region - Protein Information**ING1 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ING1 antibody - C-terminal region - Images



WB Suggested Anti-ING1 Antibody Titration: 2.5µg/ml

Positive Control: HepG2 cell lysate

There is BioGPS gene expression data showing that ING1 is expressed in HepG2



Lanes: 1: 30ug HeLa lysate, 2: 30ug HFF lysate, 3: 30ug U2OS lysate

Primary Antibody Dilution: 1:1000

Secondary Antibody: Anti-rabbit HRP

Secondary Antibody Dilution: 1:5000

Gene Name: ING1

Submitted by: Dr. Hyung Suk Oh, Knipe Lab, Harvard Medical School