

STK31 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13212**Specification**

STK31 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	B4DZ06
Other Accession	NM_032944 , NP_116562
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Predicted	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	113kDa KDa

STK31 antibody - N-terminal region - Additional Information

Alias Symbol **FLJ16102, TDRD8, SGK396**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-STK31 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


STK31 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

STK31 antibody - N-terminal region - Protein Information**STK31 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK31 antibody - N-terminal region - Images



WB Suggested Anti-STK31 Antibody Titration: 0.2-1 μ g/ml
Positive Control: HepG2 cell lysate