

Myocd antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13233**Specification**

Myocd antibody - N-terminal region - Product Information

Application	WB
Primary Accession	O6W8X1
Other Accession	NM_145136 , NP_660118
Reactivity	Human, Mouse, Rat, Rabbit, Horse, Bovine, Guinea Pig, Dog
Predicted	Mouse, Rat, Pig, Chicken, Horse, Bovine, Guinea Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	106kDa KDa

Myocd antibody - N-terminal region - Additional Information

Alias Symbol **BSAC2A, Srfcp, Myocd**

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Myocd antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions


Myocd antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

Myocd antibody - N-terminal region - Protein Information**Myocd antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Myocd antibody - N-terminal region - Images



WB Suggested Anti-Myocd Antibody Titration: 1.0 μ g/ml
Positive Control: Mouse Pancreas