

SPATA12 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13241**Specification**

SPATA12 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	Q7Z6I5
Other Accession	NM_181727 , NP_859078
Reactivity	Human
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20kDa KDa

SPATA12 antibody - N-terminal region - Additional Information**Gene ID** 353324**Alias Symbol** SRG5**Other Names**

Spermatogenesis-associated protein 12, Spermatogenesis-related protein 5, SPATA12, SRG5

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-SPATA12 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

SPATA12 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

SPATA12 antibody - N-terminal region - Protein Information**Name** SPATA12**Synonyms** SRG5**Tissue Location**


Expressed in testis.

SPATA12 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPATA12 antibody - N-terminal region - Images



WB Suggested Anti-SPATA12 Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate

SPATA12 antibody - N-terminal region - References

- Li D.,et al.Chin. Med. J. 117:1791-1796(2004).
Li D.,et al.Yi Chuan Xue Bao 31:545-551(2004).