

UBTFL1 antibody - N-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13264**Specification**

UBTFL1 antibody - N-terminal region - Product Information

Application	WB
Primary Accession	P0CB47
Other Accession	XM_926094 , XP_931187
Reactivity	Human, Rat, Guinea Pig
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46kDa KDa

UBTFL1 antibody - N-terminal region - Additional Information**Gene ID** 642623**Alias Symbol** HMGPI, C11orf27**Other Names**

Putative upstream-binding factor 1-like protein 1, UBTFL1, C11orf27

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-UBTFL1 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

UBTFL1 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

UBTFL1 antibody - N-terminal region - Protein Information**Name** UBTFL1 ([HGNC:14533](#))**Synonyms** C11orf27**Function**

Essential for proliferation of the inner cell mass and trophectodermal cells in peri-implantation development.


Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q3USZ2}. Nucleus {ECO:0000250|UniProtKB:Q3USZ2, ECO:0000255|PROSITE-ProRule:PRU00267} Note=Mainly cytoplasmic from the 4-cell stage to the morula stage Becomes nuclear at the blastocyst stage {ECO:0000250|UniProtKB:Q3USZ2}

UBTFL1 antibody - N-terminal region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UBTFL1 antibody - N-terminal region - Images

WB Suggested Anti-UBTFL1 Antibody Titration: 0.2-1 μ g/ml
Positive Control: MCF7 cell lysate