

**CROCCL2 antibody - N-terminal region**  
**Rabbit Polyclonal Antibody**  
**Catalog # AI13323****Specification**

---

**CROCCL2 antibody - N-terminal region - Product Information**

|                   |                                                         |
|-------------------|---------------------------------------------------------|
| Application       | WB                                                      |
| Primary Accession | <a href="#">Q8IVE0</a>                                  |
| Other Accession   | <a href="#">XM_057040</a> , <a href="#">XP_057040</a>   |
| Reactivity        | Human, Mouse, Rat, Rabbit, Pig, Bovine, Guinea Pig, Dog |
| Predicted Host    | Human, Rat, Bovine                                      |
| Clonality         | Rabbit                                                  |
| Calculated MW     | Polyclonal<br>47kDa kDa                                 |

**CROCCL2 antibody - N-terminal region - Additional Information**

|              |                                         |
|--------------|-----------------------------------------|
| Alias Symbol | KIAA0445, KIAA1922, dJ798A10.2, CROCCL2 |
|--------------|-----------------------------------------|

**Other Names**

Putative ciliary rootlet coiled-coil protein-like 2 protein, Ciliary rootlet coiled-coil protein pseudogene 3, CROCCP3, CROCCL2, KIAA1922

**Format**

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

**Reconstitution & Storage**

Add 50 ul of distilled water. Final anti-CROCCL2 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

**Precautions**

CROCCL2 antibody - N-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.

**CROCCL2 antibody - N-terminal region - Protein Information**

**Name** CROCCP3

**Synonyms** CROCCL2, KIAA1922


**CROCCL2 antibody - N-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **CROCCL2 antibody - N-terminal region - Images**



WB Suggested Anti-CROCCL2 Antibody Titration: 0.2-1 µg/ml  
Positive Control: MCF7 cell lysate

#### **CROCCL2 antibody - N-terminal region - References**

Nagase T., et al. DNA Res. 8:179-187(2001).  
Gregory S.G., et al. Nature 441:315-321(2006).  
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.