

C4orf33 antibody - C-terminal region
Rabbit Polyclonal Antibody
Catalog # AI13343**Specification**

C4orf33 antibody - C-terminal region - Product Information

Application	WB
Primary Accession	Q8N1A6
Other Accession	NM_001099783 , NP_001093253
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Bovine, Guinea Pig, Dog
Predicted Host	Human, Chicken, Dog
Clonality	Rabbit
Calculated MW	Polyclonal 23kDa KDa

C4orf33 antibody - C-terminal region - Additional Information**Gene ID** 132321

Alias Symbol	FLJ33703
Other Names	
UPF0462 protein C4orf33, C4orf33	

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-C4orf33 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

C4orf33 antibody - C-terminal region is for research use only and not for use in diagnostic or therapeutic procedures.


C4orf33 antibody - C-terminal region - Protein Information**Name** C4orf33**C4orf33 antibody - C-terminal region - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C4orf33 antibody - C-terminal region - Images



WB Suggested Anti-C4orf33 Antibody Titration: 0.2-1 µg/ml
Positive Control: Jurkat cell lysate

C4orf33 antibody - C-terminal region - References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Bechtel S., et al. BMC Genomics 8:399-399(2007).
Hillier L.W., et al. Nature 434:724-731(2005).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).