

Wdr92 antibody - middle region
Rabbit Polyclonal Antibody
Catalog # AI13439**Specification**

Wdr92 antibody - middle region - Product Information

Application	WB
Primary Accession	Q8BGF3
Other Accession	NM_178909 , NP_849240
Reactivity	Human, Mouse, Rat, Rabbit, Pig, Horse, Yeast, Bovine, Guinea Pig, Dog
Predicted Host	Human, Mouse, Chicken, Horse, Bovine
Clonality	Rabbit
Calculated MW	Polyclonal 39kDa KDa

Wdr92 antibody - middle region - Additional Information**Gene ID** 103784**Alias Symbol** AI553587, HZGJ, RP23-32H19.3**Other Names**

WD repeat-containing protein 92, WD repeat-containing protein Monad, Wdr92

Format

Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.

Reconstitution & Storage

Add 50 ul of distilled water. Final anti-Wdr92 antibody concentration is 1 mg/ml in PBS buffer with 2% sucrose. For longer periods of storage, store at 20°C. Avoid repeat freeze-thaw cycles.

Precautions

Wdr92 antibody - middle region is for research use only and not for use in diagnostic or therapeutic procedures.

Wdr92 antibody - middle region - Protein Information**Name** Dnaaf10 {ECO:0000312|MGI:MGI:2144224}**Function**

Key assembly factor specifically required for the stability of axonemal dynein heavy chains in cytoplasm.

Cellular Location


Dynein axonemal particle {ECO:0000250|UniProtKB:A8J3F6}

Wdr92 antibody - middle region - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Wdr92 antibody - middle region - Images



WB Suggested Anti-Wdr92 Antibody Titration: 1.0 µg/ml
Positive Control: Mouse Small Intestine

Wdr92 antibody - middle region - References

Carninci P., et al. Science 309:1559-1563(2005).
Church D.M., et al. PLoS Biol. 7:E1000112-E1000112(2009).